



Dr Aaron Chidgey

B.AppSci (Optom) (Hons) MBBS FRANZCO

Dr Aaron Chidgey brings a thoughtful, hands-on approach to eye care. He is passionate about helping patients navigate complex eye conditions with care and clarity. Dr Chidgey has special interests in medical retina (diabetic eye disease, macular degeneration and vein occlusions) and vitreoretinal surgery including trauma, complex cataract surgery (secondary lens implantation) and retinal detachments.



Dr Michael Hogden

BSc MBBS (Hons) FRANZCO DAO

Dr Michael Hogden is a retinal specialist with subspecialty training in the medical and surgical management of disorders of the vitreous, retina and macula.

Do you have a question or concern about your eye health?

To discuss your condition with an experienced ophthalmologist or optometrist, please contact The Eye Health Centre.



Brisbane

Level 11, 87 Wickham Terrace
Brisbane Qld 4000 | P 07 3831 8606

Aspley

7/1344 Gympie Road
Aspley Qld 4034 | P 07 3863 1097

Wynnum

93 Clara Street
Wynnum Qld 4178 | P 07 3026 2600


Springfield

2/95 Southern Cross Circuit
Springfield QLD 4300 | P 07 3282 4396

theeyehealthcentre.com.au

Macular Degeneration

Your guide to macular degeneration causes, symptoms and treatments.




theeyehealthcentre.com.au

What is age-related macular degeneration?

Age-related macular degeneration (AMD) is an eye condition that leads to progressive vision loss. It is a major cause of vision impairment in Australia. About one in seven Australians aged over 50 years has signs of AMD¹.

The macula is the central part of the retina, and is responsible for sharp central vision. AMD affects this area and degrades vision.



What are the symptoms of macular degeneration?

- Loss of central vision over time
- Blurry, wavy or dark spot in the central vision
- Peripheral vision remains intact
- Impact on one or both eyes.

You may not have any symptoms at all in the early stages of AMD. But as it progresses, you may experience distortion or wavy lines in your central vision.

This can progress to central vision loss.

Types of macular degeneration

There are two types of AMD: Dry and Wet. In Dry AMD there is a build up of pale yellow waste material called drusen under the macula. These can impact vision and can cause macular cells to die (atrophy) in areas and cause central vision changes. In Wet AMD, symptoms can be a more sudden onset if left untreated and can lead to central vision loss much faster. It is caused by blood vessels that grow under the macula and leak fluid or blood. This can lead to a macular scar forming, which impairs central vision.

What causes macular degeneration?

There are a number of known risk factors for macular degeneration.

- Age
- Smoking
- Obesity
- Lack of regular eye examinations
- Family history
- High blood pressure
- Poor diet, low in antioxidants

How we treat macular degeneration

There is no cure for macular degeneration but there are ways that have been shown to slow its progression.

Dry AMD

- Minimise risk factors such as high blood pressure, obesity and smoking
- Eat a diet rich in antioxidants such as dark-leafy vegetables, yellow fruits and vegetables and oily fish
- Take special vitamin tablets (Macu-Vision or Macu-Vision Plus) if advised
- Monitor your vision with an Amsler Grid regularly

Wet AMD

There are medical treatments for Wet AMD that can reduce its progression and protect further central vision loss. Some can improve vision for some people.

Intravitreal Injections

These anti-VEGF injections act to remove fluid in the retinal layers and help stop the unhealthy blood vessel growth under the macula.

There are currently 4 available anti-VEGF drugs:

- Lucentis (ranibizumab)
- Eylea (aflibercept)
- Avastin (bevacizumab)
- Beovu (brolucizumab)

The injections initially start monthly and can progress to about quarterly once the fluid or blood vessel leakage has stopped. While the goal is to stop injections without Wet AMD coming back, most patients need maintenance injections for long-term control.

Photodynamic Therapy

Since the development of intravitreal injections, photodynamic therapy is only used rarely.

It involves injecting a light activated drug called Verteporin into the bloodstream. The macula is then treated with a "cold laser" to activate Verteporin, which works to seal off abnormal blood vessels and resolve retinal fluid.

Laser Treatment

Laser treatments are uncommon in contemporary practice. They can be used for lesions that are leaking or bleeding and are located in a safe position.

Before photodynamic therapy, lasers were commonly used to treat AMD.

¹ www.mdfoundation.com.au/about-macular-disease/age-related-macular-degeneration/amd-overview